



Phytochemical screening and bioactivity of some selected Nigerian medicinal tree species grown in yankari games reserve of Bauchi state

Samaila YD Jeji^{1*}, Magnus Ezihe I², Hafiz L Zailani³

¹⁻³ Department of Science Laboratory Technology, Abubakar Tatari Ali Polytechnic, School of Science and Technology, P.O.BOX 0094, Bauchi, Nigeria

Abstract

Traditional herbal medicines play an important role in the health maintenance in developing countries, and higher plants continue to be promising sources of new medicines. In this work, we have extracted and screened six Nigerian medicinal plants for biological activity. Chloroform extracts from the aerial parts of *Aristolochia ringens*, the bark of *Chrysophyllum albidum*, the stems of *Costus afer*, the leaves of *Opilia celtidifolia*, the bark of *Terminalia catappa*, and the bark of *Vitellaria paradoxa*, were obtained and screened for phytochemical constituents, antibacterial, antifungal, and cytotoxic activities. *Aristolochia ringens* showed excellent antibacterial activity against *Bacillus cereus* and good antifungal activity against *Candida albicans*. *A. ringens* was also cytotoxic to Hep-G2 and MCF-7 cells. *Chrysophyllum albidum* showed antibacterial activity against *Escherichia coli* and cytotoxic activity to Hep-G2 cells. *Costus afer* showed good cytotoxic activity against Hep-G2 cells and antibacterial activity against *B. cereus*. *Opilia celtidifolia* extract was cytotoxic to Hep-G2 and antibacterial to *E. coli*. Neither *T. catappa* nor *V. paradoxa* extracts showed activity in the bioactivity screens in this study. The bioactivities shown in this study underscore the importance of traditional herbal medicines.

Keywords: herbal medicines, cytotoxicity, antibacterial, antifungal

Introduction

Medicinal plants constitute one of the most promising sources to obtain biologically active compounds which have been used for treatment of many human and animal diseases in various communities. Hence, natural products derived from herbs have continued to play a very important role in all divisions of human population either directly as folk medicines or indirectly in the preparation of recent drugs used as antioxidant, antibacterial, antiparasitic, antifungal, cytotoxic, anti-*Fusarium* activities for food safety and consumption [1-7]. *Aristolochia ringens* Vahl, an aromatic member of the Aristolochiaceae is known to originate from Brazil, Central America and the Caribbean [8]. The plant is commonly known by the Yoruba in the south-western part of Nigeria as 'akogun'. It is a rhizomatous aromatic climbing shrub. *Aristolochia* species are known to contain alkaloids and aristolochic acid [9]. Various *Aristolochia* species have been reported in herbal medicines since antiquity in obstetrics and in treatment of snakebite (e.g., *Echis carinatus*, saw-scaled viper) [10], festering wounds, and tumors, and they are still in use majorly in Chinese herbal medicine [11, 12]. *A. ringens* is used to treat cholera, fever, bowel troubles, ulcers, leprosy, and poisonous bites [13, 14]. The leaf, stem, and floral essential oil compositions of *A. ringens* have been previously reported [15]. Sesquiterpenoids were the main components in the leaf essential oil including β -caryophyllene (11.4%), *trans*-4(14), 5-muroladiene (13.0%), bicyclogermacrene (12.8%), spathulenol (8.0%), as well as the diterpenoid methyl copalate (10.3%). *Chrysophyllum albidum* G. Don., commonly known as white

star apple, is a forest fruit tree and belongs to the Sapotaceae family with about 800 species [16]. It is widely distributed in west and central tropical Africa [17-19]. In Nigeria, it is found along forest savannah transitional and coastal savannah zones. The plant often grows to a height of 25-37 m in height with a mature girth varying from 1.5 to 2 m [20]. The tree has dark green, pale tawny with silver-white when mature, flowers appear on the leaf axils and fruit spherical slightly pointed at the tip with about 3.2 cm in diameter, greenish-grey when unripe, turning orange-red when matured. Locally the plant is named 'agbalumo', (Yoruba) 'udara' (Igbo) according to the specific area. *C. albidum* has been used in traditional herbal medicine to treat malaria and yellow fever, while the leaf is used as an emollient and for the treatment of skin eruptions, stomach ache and diarrhea [21]. Earlier investigators have reported that extracts of the seeds and roots of *C. albidum* have demonstrated good antimicrobial (e.g., *Escherichia coli*), anti-inflammatory, anti-diarrheal and antihemorrhoidal properties [22]. *Costus* (Ginger lily) is a genus of stout, perennial and rhizomatous herbs of the Costaceae comprised of 150 species found in tropical belt of West African countries including Cameroon and South Africa [23-25]. *Costus afer* Ker Gawl is an erect herbaceous plant growing up to 4 m with white and yellow flowers [26]. The succulent stem and root contain several bioactive metabolites as a remedy for headache, fever, rheumatism, cough, sleeping sickness and stomach ache [27]. *C. afer*, locally called bush cane, 'ireke omode' or 'teteregun' (Yoruba) and 'opete' (Igbo), is used in West African traditional medicine for ailments such as laxative, hepatic

disorders and miscarriage. The plant has diverse pharmacological properties viz. anti-inflammatory, rheumatism, arthritis, helminthic, epileptic attack, hemorrhoids, as diuretics, and also had served as an antidote for food poisoning [28, 29]. Laboratory study of the stem of *C. afer* had shown that the plant possessed potent antioxidant activity [30]. Several groups have investigated the essential oil [29], topical anti-inflammatory activity [30] as well as *in vitro* and *in vivo* pharmacological activities of the methanol leaf extract [28].

Opilia celidifolia (Guill. & Perr.) Endl. ex Walp. (Opiliaceae) is a spreading liana, heavily-branched shrub or tree up to a height of 10 m, common in savannah or fringing forest. It is prevalent in western Africa from Nigeria to Senegal and disseminated over the arid parts of tropical Africa [24]. Medicinally, the plant decoction is taken as a remedy for wound healing, malaria fever, abdominal pains, internal worms and as an appetite-enhancing agent [31-36]. The plant, locally called 'koyinsorun', is used in West African traditional medicine for treating sleeping sickness and leprosy [24]. *Terminalia catappa* (tropical almond) are trees or shrubs in the Combretaceae family. The genus is made up of nearly 200 species [9] and is native to tropics and sub tropics regions of Africa, India and Southeast Asia including Thailand and Malaysia. *T. catappa* is a tree, up to 35 m in height with an upright, symmetrical crown and produces fruits 5-10 cm long, broad, green at first, then yellow and finally red when ripe with a flesh surrounding a large fibrous nuts. All parts of the plant had been studied extensively. Ethnobotanical, the aqueous bark extract is used traditionally for treatments of wounds, boils, dysentery and diarrhea. Leaf decoctions of *T. catappa* have been broadly used in local traditional medicine for the treatment of liver ailments, headache, colic, and as a cardiogenic and diuretic. Petroleum ether, methanol, and aqueous extracts of *T. catappa* have also demonstrated antidiabetic activity.

Vitellaria paradoxa C.F. Gaertn. [syn. *Butyrospermum paradoxum* (C.F. Gaertn.) Hepper], belonging to the family Sapotaceae, is a small to medium-sized tree, 10-15 m high, much branched, dense, spreading, round to hemispherical crown. The tree grows naturally in the wild of the dry savannah belt of West Africa, from Senegal in the west to Sudan in the east and onto the foothills of the Ethiopian Mountains. It is rich in oil and serves as a source of edible oil for many households in the northern part of Nigeria. *V. paradoxa* has many applications in folkloric medicine. It is commonly called 'shea butter' (English) and 'emi-emi' (Yoruba) in Nigeria. Shea butter contains high levels of UV absorbing triterpene cinnamates, tocopherols (vitamin E), and phytosterols. Baidu confirmed that shea butter contains high anti-oxidant properties and (5-15%) of unsaponifiable lipids, which include phytosterols such as campesterol, stigmaterol, β -sitosterol and α -spinosterol, as well as triterpenes such as cinnamic acid esters of α - and β -amyrin, parkeol, butyrospermol, and lupeol. Its anti-oxidant properties have led to its use to protect the skin from sunburn, eczema and as a skin rejuvenator. Additionally, shea kernels contain about 0.4% catechin polyphenolics such as gallic acid, catechin, epicatechin, epicatechin gallate, gallocatechin, epigallocatechin, epigallocatechin gallate, as well as quercetin

and *trans*-cinnamic acid. A phytochemical screen of the methanol bark extract of *V. paradoxa* has revealed the presence of carbohydrates, saponins, steroids, and alkaloids.

Materials and Methods

Plant Materials

The aerial parts of *A. ringens* and the stem bark of *C. albidum* were procured from a local market at Idi Oro market, Lagos State (6.56° N, 3.3° E); the mature stems of *C. afer* (tropical almond) were procured from a local market at Lagos Island, Lagos State; mature leaves of *O. celidifolia* and stem bark of *V. paradoxa*. (shea-butter tree) were procured from a local market at Agege, Lagos State; the stem bark of *T. catappa* was collected from the premises of Lagos State University central mosque. All plant materials were identified at the Herbarium unit of the Department of Botany, University of Lagos. The plant materials for each plant were dried in air for five days, pulverized using a grinding machine, and stored in polyethylene bags. For each plant, 750 g of plant material was soaked in 1.5 L chloroform for 24 h. Each crude extract was filtered using Whatman filter paper No 42 (125 mm). The filtrates were concentrated under reduced pressure at 80°C using a rotary evaporator and stored under refrigeration at 4°C. The extract yields of each extract were determined and are represented as percentages based on dry mass of the plant materials. All the solvents, chemicals, and reagents used were of analytical grade and were purchased either from Tunnex Chemicals, Alapere (Lagos) or from Olaolu Chemicals, Igbo Elerin, (Lagos).

Qualitative Phytochemical Screening

Phytochemical screening was performed on the extracts using standard protocols to identify chemical constituents.

Screening for Alkaloids: The crude extract (0.5 g) was mixed with 5 mL of 1% HCl on a steam bath and filtered while hot. The residue was rinsed with distilled water and 1 mL of the filtrate was treated with a few drops of Wagner's reagent. A positive alkaloid test was revealed by a reddish brown precipitate.

Screening for Flavonoids: The extract (0.2 g) was dissolved in water (2 mL) and filtered. Aqueous sodium hydroxide (2 mL, 10%) was added. The appearance of a yellow color indicated the presence of flavonoids.

Screening for Phenolics: The chloroform extract (0.2 g) was diluted with 50% ethanol and 3 drops of 10% ferric chloride were added. A deep bluish-green solution indicated of the presence of phenols.

Screening for Tannins: A portion of the extract was dissolved in water, after which the solution was clarified by filtration. A 10% ferric chloride solution was added to the filtrate. The appearance of a bluish-black color indicated a positive test for tannins.

Screening for Anthraquinones: Crude extract (0.5 g) was shaken with 10 mL of benzene and filtered. A 10%. Journal of Pharmacognosy and Phytochemistry aqueous ammonia solution was added to the filtrate and the mixture was shaken.

The formation of a pink, red or violet color in the aqueous phase indicates the presence of anthraquinones.

Screening for Cardiac Glycosides: Crude extract (0.5 g) was dissolved in glacial acetic acid (2 mL) containing 1 drop of 10% ferric chloride solution. The solution was under layered with 2 mL of concentrated sulfuric acid. Formation of a brown ring at the phase interface indicated the presence of deoxy sugars, characteristic of cardiac glycosides.

Screening for Phlobatannins: Crude extract (0.2 g) was mixed with 10 mL water. A few drops of 1% HCl were added and the mixture boiled. Formation of a red precipitate indicated the presence of phlobatannins.

Screening for Terpenoids: Acetic anhydride (0.5 mL) was mixed with 0.2 g extract dissolved in 50% ethanol. A few drops of concentrated sulfuric acid were added. Formation of a bluish green precipitate indicated the presence of terpenoids.

Screening for Steroids: Crude extract (0.2 g) was dissolved in 2 mL of glacial acetic acid and the solution heated, allowed to cool, and filtered. The residue was dissolved in 0.5 mL of acetic anhydride followed by the addition of few drops of concentrated sulfuric acid. A color change from violet to blue or bluish-green was a positive indication for the presence of steroids.

Quantitative Phytochemical Screening

Determination of Total Alkaloids: The crude extract (5 g) was weighed into a 250-mL beaker and 200 mL of 10% acetic acid in ethanol was added and covered and allowed to stand for 4 h. This was filtered and the extract was concentrated on a water bath to one-quarter of the original volume. Concentrated aqueous ammonia was added drop wise to the extract until the precipitation was complete. The whole solution was allowed to settle and the precipitated was collected and washed with dilute ammonium hydroxide and then filtered. The solid alkaloid residue was dried and weighed.

Determination of Total Phenolics, Phenanthroline Method: The chloroform extract (500 mg) was dissolved in methanol, 1 mL of 0.2% FeCl₃ solution in methanol, and 0.5 mL of 0.5% 1,10-phenanthroline solution in methanol were placed into a 10-mL volumetric flask and diluted to a final volume of 10 mL with methanol. The solution was mixed and left at room temperature in the dark. After 20 min, the absorbance of an orange-red solution was measured at 510 nm against a reagent blank (1 mL of 0.2% FeCl₃ and 0.5 mL of 0.5% 1,10-phenanthroline, diluted to 10 mL). The results are expressed as mg of gallic acid equivalents per 100 g plant extract.

Determination of Total Phenolics, Folin-Ciocalteu Method: The total phenolic content (TPC) of the chloroform extracts was determined using the Folin-Ciocalteu method. Each extract (100 mg) was dissolved in 2 mL acetone to which was then added 2 mL of 50% Folin-Ciocalteu reagent [64], followed by addition of 5 mL of 20% Na₂CO₃ solution. The mixture was shaken vigorously and diluted to 10 mL.

After 30 min the absorbance was measured at 730 nm. The total phenolic content is expressed as mg of gallic acid equivalents per 100 g plant extract.

Determination of Total Flavonoids: Crude extract (10 g) was exhaustively extracted with 100 mL of 80% aqueous methanol at room temperature. The solution was filtered and the filtrate transferred into a crucible and evaporated to dryness over a water bath. Total flavonoid content is expressed as the mass of the residue after drying.

Antimicrobial Screening: The chloroform extracts were screened for antibacterial activity against Gram-positive bacteria, *Bacillus cereus*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*, and Gram-negative bacteria, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Serratia marcescens*. Minimum inhibitory concentrations (MIC) were determined using the microbroth dilution technique. Dilutions of the chloroform extracts were prepared in cation-adjusted Mueller Hinton broth (CAMHB) beginning with 50 μ L of 1% w/w solutions of each crude extract in DMSO plus 50 μ L CAMHB. The extracts were then serially diluted (1:1) in CAMHB in 96-well plates. Microorganisms at a concentration of approximately 1.5×10^8 colony forming units (CFU)/mL were added to each well. Plates were incubated at 37°C for 24 h; the lowest concentration without turbidity was defined as the final minimum inhibitory concentration (MIC). DMSO was used as a negative control and Gentamicin was used as a positive antibiotic control. Antifungal activity was determined as described above using *Candida albicans* in yeast-mold (YM) broth with approximately 7.5×10^7 CFU/mL. Antifungal activity against *Aspergillus niger* and *Botrytis cinerea* were determined as above using potato dextrose broth inoculated with *A. niger* hyphal culture and grey mold horticulture, respectively, diluted to a McFarland turbidity of 1.0. Amphotericin B was the positive control.

Cytotoxicity Screening

Human hepatocarcinoma cell line (Hep-G2, CRL-11997™) cells were grown in complete medium (DMEM: F12 containing L-glutamine and sodium bicarbonate, 10% FBS, and 1% penicillin/streptomycin) incubated at 37°C in a 5% CO₂ environment. Once 80 - 90% confluent, the cells were washed with phosphate buffered saline (PBS), treated with 0.25% (w/v) of Trypsin/EDTA, counted and suspended in fresh complete media. About 5×10^5 cells/well (100 μ L) were seeded into 96 well plates and incubated for 24 hours to attach. Cells were then treated with the extracts at a final concentration of 10 and 50 μ g/mL in 200 μ L medium for 72 hours. Thereafter, the medium was removed, and DMEM: F12 medium containing MTT (5 mg/mL in PBS) was added to the cells and incubated for 1 h. The MTT-containing medium was then carefully removed and replaced with DMSO (200 μ L per well), the plate was then gently mixed to dissolve the formazan crystals. Absorbance was measured at 550 nm. All extracts were tested in triplicate. MCF-7 (human breast tumor, ATCC No. HTB-22) cells were grown in RPMI 1640 supplemented with 10% Fetal bovine serum (FBS), 30 mM HEPES, sodium bicarbonate, and 100,000 units penicillin/streptomycin (10 mg/L) at pH=7.35. MCF-7 cells

were plated into 96-well cell culture plates at a concentration of 1.2×10^4 cells/well and a volume of 100 μ L in each well and incubated at 37°C and 5% CO₂ for 48 hours. After 48 hours, the cells reached 70-80% confluent growth. The supernatant fluid was carefully aspirated and replaced with 100 μ L growth medium containing 1 μ L of extract (1% in DMSO), giving a final concentration of 100 μ g/mL (100 ppm). Assays were repeated at 10 μ g/mL (10 ppm). The plate was then incubated at 37°C and 5% CO₂ for 48 hours, after which the supernatant liquid was gently aspirated from each well. Into each well, 100 μ L of MTT solution was added and the pre-read absorbance was immediately measured spectrophotometrically at 570 nm (using a Molecular Devices Journal of Pharmacognosy and Phytochemistry SpectraMax Plus 384 microplate reader). The plate was incubated at 37°C and 5% CO₂ for 4 h, after which the supernatant liquid was removed and DMSO (100 μ L) was used to dissolve the purple

formazan crystals. The amount of formazan produced was determined spectrophotometrically at 570 nm. DMSO, and tingenone (100 μ g/mL) served as negative and positive controls, respectively. Solutions were added to wells in eight replicates. Average absorbances, standard deviations, and percent kill ratios (% killcompound / % killcontrol) were calculated. Median inhibitory concentrations (IC₅₀) were determined using the Reed-Muench method.

Results and Discussion

Extraction

The chloroform extraction yields and descriptions of the Medicinal tree plants in Bauchi State are summarized in Table 1. Bark extractions with chloroform gave larger yields than leaves or herbaceous plant parts, which suggests larger concentrations of relatively non-polar extractables in the barks.

Table 1: Nature and yield of crude chloroform extracts from medicinal trees plants.

Plant Extract	Extract color	Extract texture	Extract yield
<i>Aristolochia ringens</i> aerial parts	Dark green	Semi-solid	65.00g
<i>Chrysophyllum albidum</i> bark	Brown	Semi-sticky powder	100.00g
<i>Opilia celtidifolia</i> leaves	Brown	Solid powder	57.00g
<i>Terminalia catappa</i> bark	Brown	Solid powder	60.00g
<i>Vitellaria paradoxa</i> bark	Dark brown	Semi-solid	61.00g

Plant Extract Extract color Extract texture Extract yield

Aristolochia ringens aerial parts Dark brown Sticky semi-solid powder 84.00 g (7.0%) *Chrysophyllum albidum* bark Dark green Semi-sticky powder 120.00 g (10.0%) *Costus afer* stems Brown Solid powder 75.32 g (5.94%) *Opilia celtidifolia* leaves Brown Solid powder 70.00 g (5.84%) *Terminalia catappa* bark Dark brown Semi-solid powder 153.55 g (14.00%) *Vitellaria paradoxa* bark Dark brown Semi-solid powder 150.00 g (13.0%)

Phytochemical Screening

Qualitative phytochemical screening was carried out on the crude chloroform extracts, and the results are summarized in Table 2. Quantitative assessment of phytochemical constituents is summarized. Not surprisingly, polar phytochemicals such as phenolics and tannins were found in only two crude chloroform extracts. Conversely, alkaloids were found in five of the six extracts, while flavonoids and terpenoids were found in four extracts.

Antimicrobial Screening

Each of the crude chloroform extracts was screened for antimicrobial activity against a panel of Gram-positive bacteria (*Bacillus cereus*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*), Gram-negative bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*, and *Serratia marcescens*), and fungi (*Aspergillus niger*, *Botrytis cinerea*, and *Candida albicans*). Antimicrobial activities as minimum inhibitory concentrations are summarized. The most susceptible organisms in this study were *B. cereus* and *E. coli* with two extracts showing MIC < 100 μ g/mL and one extract with MIC = 156 μ g/mL. *Staphylococcus epidermidis* and *Serratia marcescens* were particularly insensitive to the plant extracts. Likewise, neither *A. niger* nor *B. cinerea* were susceptible to the Nigerian extracts. *Candida albicans*, however, was somewhat sensitive to *A. ringens* extract (MIC = 156 μ g/mL).

Table 2: Qualitative analysis of phytochemical constituents of medicinal tree plants.

Plant Extracts	Phenolics	Tannins	Phlobatannins	Alkaloids	Flavanoids	Cardiac glycosides	Steroids	Terpenoids	Antraquinones
<i>Aristolochia ringens</i> aerial parts	-	-	-	+++	++	-	+	-	+
<i>Chrysophyllum albidum</i> bark	-	-	-	+++	++	-	+	-	+
<i>Opilia celtidifolia</i> leaves	++	++	++	+++	++	-	-	-	+
<i>Terminalia catappa</i> bark	+	++	+	+++	+	-	+	+	-
<i>Vitellaria paradoxa</i> bark	-	-	+	+++	-	-	+	-	-

Plant Extract Phenolics Tannins Phlobatannins Alkaloids Flavonoids Cardiac glycosides Steroids Terpenoids Anthraquinones

Plant Extract Phenolics (mg/100 g) 1,10-phenanthroline method Phenolics (mg/100 g) Folin-Ciocalteu method Alkaloids (mg/100 g) Flavonoids (mg/100 g) Cardiac glycosides extracts.

Plant Extract Gram-positive bacteria Gram-negative bacteria Fungi

B. cereus *S. aureus* *S. epidermidis* *E. coli* *P. aeruginosa* *S. marcescens* *A. niger* *B. cinerea* *C. albicans*

Aristolochia ringens aerial parts < 19.5 > 2500 > 2500 > 2500 625 1250 625 625 156 *Chrysophyllum albidum* bark 313 2500 > 2500 < 19.5 313 > 2500 625 625 625 *Costus afer* stems 78 313 > 2500 156 313 > 2500 1250 625 > 2500 *Opilia celtidifolia* leaves 156 625 > 2500 78 156 > 2500 1250 625 > 2500 *Terminalia catappa* bark > 2500 1250 > 2500 2500 313 > 2500 > 2500 625 > 2500 *Vitellaria paradoxa* bark > 2500 1250 > 2500 2500 313 > 2500 > 2500 625 > 2500

Cytotoxicity Screening

The crude chloroform extracts were screened for *in-vitro* cytotoxic activity against Hep-G2 human hepatocellular carcinoma and MCF-7 human breast adenocarcinoma cells.

Plant Extract Hep-G2 MCF-7

Aristolochia ringens aerial parts 16.4 ± 1.8 81.6 ± 12.8

Chrysophyllum albidum bark 23.1 ± 4.8 > 100

Costus afer stems 17.6 ± 1.3 > 100

Opilia celtidifolia leaves 24.7 ± 6.8 > 100

Terminalia catappa bark > 50 > 100

Vitellaria paradoxa bark > 50 129 ± 14

Aristolochia ringens

Phytochemical screening of *A. ringens* chloroform extract revealed the presence of alkaloids, flavonoids, steroids, and anthraquinones. This extract showed excellent antibacterial activity against *B. cereus* (MIC < 19.5 µg/mL) and antifungal activity against *C. albicans* (MIC = 156 µg/mL). *A. ringens* extract was also cytotoxic to both Hep-G2 and MCF-7 cell lines (IC₅₀ = 16.4 and 81.6 µg/mL, respectively). Previous studies on *A. ringens* have shown that the root extracts of *A. ringens* have exhibited insecticidal activity as well as broad spectrum cytotoxic activity, while the methanol extract of the plant showed *in-vivo* antitrypanosomal activity. The volatile components from the stems and leaves of *A. ringens* have been identified [15]. The stems were dominated by the monoterpenoids limonene (20.0%), *p*-cymene (17.8%), α -phellandrene (16.1%), and linalool (6.5%), while the leaves were rich in sesquiterpenoids β -caryophyllene (11.4%), *trans*-4(14), 5-muroladiene (13.0%), and spathulenol (8.0%), and the diterpenoid methyl copalate (10.3%). The diterpenoid *ent*-labd-8(17)-en-15-oic acid has also been identified in *A. ringens*.

Chrysophyllum albidum

The crude chloroform bark extract of *C. albidum* showed selective *in-vitro* cytotoxicity to Hep-G2 cells (IC₅₀ = 23.1 µg/mL). In addition, the extract showed excellent antibacterial activity against *E. coli* (MIC < 19.5 µg/mL). In a previous

study, the ethanol leaf extract of *C. albidum* showed antibacterial activity against Gram-positive *S. aureus* and *Streptococcus pyogenes*, but MIC values were not determined. In addition, the methanol bark extract of *C. albidum* has shown *in-vivo* antiplasmodial activity in mice, but was relatively non-toxic to the mice (LD₅₀ = 1850 mg/kg). Phytochemical screening of the chloroform bark extract of *C. albidum* showed positive results for alkaloids, flavonoids, cardiac glycosides, and steroids (Tables 2). Previous phytochemical analysis of the stem bark of *C. albidum* showed large quantities of alkaloids, flavonoids, cardiac glycosides, and saponins [22], complementing the results of this current study. Leaf extracts of *C. albidum* have also tested positive for alkaloids, flavonoids, cardiac glycosides, and steroids.

Costus afer

The chloroform extract of the stems of *C. afer* tested positive for phenolics, tannins, phlobatannins, alkaloids, flavonoids, cardiac glycosides, and terpenoids. The phytochemical screening results corroborate previous studies of the stem extract that indicated alkaloids, saponins, triterpenoids and steroids, flavonoids, phlobatannins, and tannins [25, 28]. The flavonoid glycoside kaempferol 3-*O*- α -L-rhamnopyranoside has been isolated from the aerial parts of *C. afer*. Volatile constituents of the stem have included fatty acids and fatty acid derivatives (palmitic acid, (*Z*)-vaccenic acid, oleic acid, stearic acid) as well as phytosterols (campesterol and stigmasterol). The chloroform stem extract in this study showed antibacterial activity against *B. cereus* (MIC = 78 µg/mL) and *E. coli* (MIC = 156 µg/mL), but no antifungal activity. In addition, *C. afer* extract showed good, selective cytotoxicity against Hep-G2 cells. The juice and the leaf extract of *C. afer* have shown moderate antiparasitic activity against *Entamoeba histolytica*.

Opilia celtidifolia

The chloroform leaf extract of *O. celtidifolia* showed selective *in-vitro* cytotoxicity to Hep-G2 cells (IC₅₀ = 24.7 µg/mL) and moderate antibacterial activity to *B. cereus*, *E. coli*, and *P. aeruginosa* (MIC = 156, 78, and 156 µg/mL, respectively). *O. celtidifolia* methanol extract had previously shown cytotoxic activity to HeLa and A431 cells. *Opilia celtidifolia* is used traditionally to treat malaria and the dichloromethane leaf extract has shown *in-vitro* antiplasmodial and antitrypanosomal activity. Phytochemical screening of the extract indicated the presence of alkaloids, phlobatannins, terpenoids, and anthraquinones. Leaf extracts of *O. celtidifolia* have revealed triterpenoid saponins.

Terminalia catappa

The crude chloroform bark extract of *T. catappa* showed positive phytochemical tests for polyphenolics, including tannins, phlobatannins, and flavonoids, as well as terpenoids and anthraquinones. The extract was, however, inactive in the bioactivity screens in this present study. Previous work with *T. catappa* bark extracts have shown the ethanol bark extract to be weakly antibacterial while the hexane bark extract showed antifungal activity. Minimum inhibitory concentrations were not determined in either of these previous studies, however.

Sterols, triterpenoids, and saponins have been isolated and identified in the bark extracts of *T. catappa*.

Vitellaria paradoxa

Vitellaria paradoxa chloroform bark extract tested positive for only alkaloids and terpenoids. Likewise, the bark extract was neither antimicrobial nor cytotoxic in our bioassays. In contrast, the methanol bark extract has shown *in-vitro* ~ 1435 ~ Journal of Pharmacognosy and Phytochemistry cytotoxic activity against several tumor cell lines with IC50 ranging from 24 to 67 µg/mL. Likewise, the ethanol bark extracts of *V. paradoxa* have shown marginal activity against several bacterial and fungal strains (MIC ≥ 50 mg/mL). The dichloromethane bark extract of *V. paradoxa* showed *in vitro* antiplasmodial activity. Sterols, triterpenoids, and flavonoids have been isolated and characterized from *V. paradoxa* bark.

Conclusions

The results of this investigation suggest that the use of some of these plants in traditional herbal medicines can be scientifically justified. Traditional herbal medicines play a major role in the health care in many developing nations and such herbal medicines may provide new chemotherapeutic agents for discovery and development as conventional singleentity drugs.

References

1. Antony R, Uddin SJ, Grice ID, Tiralongo E. Cytotoxic activity screening of Bangladeshi medicinal plant extracts. *Journal of Natural Medicines*. 2014; 68:246-252.
2. Alfatemi SMH, Rad JS, Rad MS, Mohsenzadeh S, da Silva JAT. Chemical composition, antioxidant activity and *in vitro* antibacterial activity of *Achillea wilhelmsii* C. Koch essential oil on methicillin-susceptible and methicillin-resistant *Staphylococcus aureus* spp. *3 Biotech*. 2015; 5:39-44.
3. Bhagwat MK, Datar AG. Antifungal activity of herbal extracts against plant pathogenic fungi. *Archives of Phytopathology and Plant Protection*. 2014; 47:959-965.
4. Horn D, Duraisingh MT. Antiparasitic chemotherapy: From genomes to mechanisms. *Annual Review of Pharmacology and Toxicology*. 2014; 54:71-94.
5. Sharifi Rad J, Hoseini-Alfatemi SM, Sharifi Rad M, Iriti M. Free radical scavenging and antioxidant activities of different parts of *Nitraria schoberi* L. *Journal of Biologically Active Products from Nature*. 2014; 4:44-51.
6. Tangarife-Castaño V, Correa-Royero JB, Roa-Linares VC, Pino-Benitez N, Betancur-Galvis LA, Duráncd DC, *et al.* Anti-dermatophyte, anti-*Fusarium* and cytotoxic activity of essential oils and plant extracts of *Piper* genus. *Journal of Essential Oil Research*. 2014; 26:221-227.
7. Znati M, Jannet HB, Cazaux S, Bouajila J. Chemical composition, biological and cytotoxic activities of plant extracts and compounds isolated from *Ferula lutea*. *Molecules*. 2014; 19:2733-2747.
8. Watson L, Dallwitz MJ. *The Families of Flowering Plants: Interactive Identification and Information Retrieval*, CSIRO, East Melbourne, Victoria, 1994.
9. Mabberley DJ. *Mabberley's Plant-Book*, 3rd Ed, Cambridge University Press, Cambridge, UK, 2008.
10. Meenatchisundaram S, Prajish, Parameswari G, Subbraj T, Michael A. Studies on antivenom activity of *Andrographis paniculata* and *Aristolochia indica* plant extracts against *Echis carinatus* venom. *Internet Journal of Toxicology*. 2008; 6:1-7.
11. Balachandran P, Wei F, Lin RC, Khan IA, Pasco DS. Structure activity relationships of aristolochic acid analogues: Toxicity in cultured renal epithelial cells. *Kidney International*. 2005; 67:1797-1805.
12. Zhou X, Zheng C, Sun J, You T. Analysis of nephrotoxic and carcinogenic aristolochic acids in *Aristolochia* plants by capillary electrophoresis with electrochemical detection at a carbon fiber microdisk electrode. *Journal of Chromatography A*. 2006; 1109:152-159.
13. Krishnaraju AV, Rao TVN, Sundararaju D, Vanisree M, Tsay HS, Subbaraju GV. Assessment of bioactivity of Indian medicinal plants using brine shrimp (*Artemia salina*) lethality assay. *International Journal of Applied Science and Engineering*. 2005; 3:125-134.
14. Kanjilal PB, Kotoky R, Couladis M. Chemical composition of the stem oil of *Aristolochia indica* L. *Journal of Essential Oil Research*. 2009; 21:24-25.
15. Stashenko EE, Ordóñez SA, Marín NA, Martínez JR. Determination of the volatile and semi-volatile secondary metabolites, and aristolochic acids in *Aristolochia ringens* Vahl. *Journal of Chromatographic Science*. 2009; 47:817-821.
16. Ehiagbonare JE, Onyibe HI, Okoegwale EE. Studies on the isolation of normal and abnormal seedlings of *Chrysophyllum albidum*: A step towards sustainable management of the taxon in the 21st century. *Scientific Research and Essays*. 2008; 3:567-570.
17. Adebayo AH, Abolaji AO, Opatá TK, Adegbenro IK. Effects of ethanolic leaf extract of *Chrysophyllum albidum* G. on biochemical and haematological parameters of albino Wistar rats. *African Journal of Biotechnology*. 2010; 9:2145-2150.
18. Amusa NA, Ashaye OA, Oladapo MO. Biodeterioration of the African star apple (*Chrysophyllum albidum*) in storage and the effect on its food value. *African Journal of Biotechnology*. 2003; 2:56-59.
19. Bada SO. Preliminary information on the ecology of *Chrysophyllum albidum* G. Don in West Africa and Central Africa. In: *Proceedings of a National workshop on the potentials of the star apple in Nigeria*; Denton DA, Ladipo DO, Adetoro MA, Sarumi MB, Eds, 1997, 16-25.
20. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. *Agroforestry Database: A tree reference and selection guide version 4.0* <http://www.worldagroforestry.org/output/agroforestrydatabase> (accessed Aug 19, 2017).
21. Idowu TO, Iwalewa EO, Aderogba MA, Akinpelu BA, Ogundaini AO. Antinociceptive, anti-inflammatory and antioxidant activities of eleagnine: An alkaloid isolated from *Chrysophyllum albidum* seed cotyledons. *Journal of Biological Sciences*. 2006; 6:1029-1034.
22. Okoli BJ, Okere OS. Antimicrobial activity of the phytochemical constituents of *Chrysophyllum albidum* G. Don Holl. (African star apple) plant. *Journal of Research in National Development*, 2010, 8.

23. Edeoga HO, Okoli BE. Chromosome numbers of *Costus lucanusianus* (Costaceae) in Nigeria. *Folia Geobotanica*. 2000; 35:315-318.
24. Burkhill HM. *Useful Plants of West Tropical Africa*, Royal Botanic Gardens, Kew, UK, 2000, 1.
25. Ukpabi CF, Agbafor KN, Ndukwe OK, Agwu A, Nwachukwu SN. Phytochemical composition of *Costus afer* extract and its alleviation of carbon tetrachloride~1436 ~ *Journal of Pharmacognosy and Phytochemistry induced hepatic oxidative stress and toxicity*. *International Journal of Modern Botany*. 2012; 2:120-126.
26. Steentoft M. *Flowering Plants in West Africa*, Cambridge University Press, Cambridge, UK, 2008.
27. Iwu MM. *Handbook of African Medicinal Plants*, 2nd Ed, CRC Press, Boca Raton, Florida, 2014.
28. Anyasor GN, Ogunwenmo KO, Oyelana OA, Akpofunure BE. Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costus afer* Ker Gawl. (Costaceae). *African Journal of Biotechnology*. 2010; 9:4880-4884.
29. Taiwo AO, Bolanle AA. The leaf essential oil of *Costus afer* Ker-Gawl from Nigeria. *Flavour and Fragrance Journal*. 2003; 18:309-311.
30. Moody JO, Okwagbe KE. Topical anti-inflammatory activity of *Costus afer*. *Nigerian Journal of Natural Products and Medicine*. 2003; 7:46-48.