

Nutraceutical properties of *Stellaria media* (L.) Chickweed

Priti Kushwaha¹, IP Kumhar²

¹ Research Scholar, Department of Botany, S.G.S. Govt. P.G. College, Sidhi, Madhya Pradesh, India

² Professor, Department of Botany, S.G.S. Govt. P.G. College, Sidhi, Madhya Pradesh, India

Abstract

Many of herbs are used as dietary source and plays a vital role in improving our health. Antioxidant compounds in diet play an important role as a health protecting factor. Various herbs have been reported to exhibit antioxidant activity which contains many different antioxidant compounds (i.e. vitamin C and E, carotenoids and phenolic compounds), they serve as free radical scavengers. The present study was subjected to investigate the nutraceutical property and antioxidant activity of *Stellaria media* L. (Family-Caryophyllaceae) found available in Sidhi district agro-Climatic Condition. The ascorbic acid and phenolic content was observed in *S. media*. Free radical scavenging activity was evaluated using 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical. The value of methanolic extract observed $1021 \pm 0.67 \mu\text{g/ml}$ in *S. media*. The study reveals that the use of this plant as therapeutic agent would exert several beneficial effects by virtue of its antioxidant activity and also can be a rich source of nutrition in our diet system.

Keywords: *Stellaria media*, antioxidant activity, DPPH, total phenolic content, ascorbic acid, gallic acid

1. Introduction

For sustaining healthy and active life, our diet should contain proper nutritional elements which have health promoting as well as health protective activity. The dietary habits of population in different region of the world have been determined mainly by the availability of foods locally and local practices. Production of certain free radicals such as super oxide, hydrogen peroxide, hydroxyl and nitric oxide radicals are the consequence of body's natural metabolic process which may be involved in the onset of many diseases such as cancer, rheumatoid arthritis, and atherosclerosis as well as in degenerative processes associated with ageing [1]. Almost all organisms are well protected against free radical damage by enzymes such as superoxide dismutase and catalase, or compounds such as ascorbic acid, tocopherols and glutathione [2]. When the mechanism of antioxidant protection becomes unbalanced by factors such as aging, deterioration of physiological functions may occur, resulting in diseases and accelerating aging. However the natural antioxidant such as vitamin C, E, carotenoids, phenolic compounds, etc. that are present in herbs and spices are responsible for inhibiting or preventing the deleterious consequences of oxidative stress exerted by the reactive oxygen species (ROS) [3]. It has been reported that the antioxidant activity of plant materials are well correlated with the content of their phenolic compounds [4-5]. Phenolic compounds, especially phenolic acids and flavonoids, are ubiquitously present in vegetables, fruits, seeds, tea, wines and juices; thus they are an integral part of the human diet. In a broad sense, nutritive foods are those foods, which not only provide nutrition but at the same time have health protective and health promoting properties. Present paper deals the adaptability to almost all environmental conditions makes common chickweed a very

widespread and successful weed. It can start producing seeds within five weeks of germination and can continue to produce seeds for weeks or even months thereafter. Seeds can germinate (sprout) soon after they disperse from the plant and require no further ripening as do seeds of many other weeds.

It is highly medicinally important weed found as chick weed in Sidhi district.



Fig 1: Common Chick weed *Stellaria media* L.

2. Material and Methods

Methods of analysis

Chemical analysis was done on moisture free basis. Analysis was carried out to estimate the macro nutrient components viz. total protein, total carbohydrates, crude fibre and ascorbic acid and antioxidant activity of the samples.

Total Protein Estimation

The total protein content of *S. media* was estimated by following the method developed by Lowry *et al.* (1962) [6]. Extraction was carried out with buffers used for the enzyme assay. 500 mg of the sample was grinded well with a mortar and pestle in 5ml of the buffer and after centrifuging; the supernatant was used for protein estimation. The reading was taken in UV-Vis Spectrophotometer at 660nm and the amount of protein present was calculated by plotting the value in a standard curve of Bovine Serum Albumin (BSA).

Total Carbohydrate Estimation

The total carbohydrate content of the samples was estimated by Anthrone method [7]. 100mg dried samples were hydrolyzed with 2.5 N HCl for about 3 hours in a boiling water bath. Sodium carbonate was added to neutralize the extracts. Subsequently the extracts were centrifuged and supernatant were collected. The residue was washed thrice with distilled water and all the supernatants were pooled and final volume was adjusted to 100ml. 0.5ml of the extracts were taken and volume made up to 1ml distilled water. 4ml of Anthrone reagent was added to the above solution. Absorbance was taken in UV-Vis spectrophotometer at 630nm and the amount of carbohydrate present was calculated by plotting the value in a standard curve of standard Glucose solution.

Ascorbic Acid Content Estimation

The amount of ascorbic acid present in the samples was calculated by extracting the sample in 4% oxalic acid and titrating the extract against the 2, 6-dichloro phenol indophenols dye until the end point where pink colour appears that persist for a few minutes [8]. The amount of dye consumed is equivalent to the amount of ascorbic acid present in the samples. Standard ascorbic acid solution was used as the reference and the calculation was done by the following formulae:

Amount of ascorbic acid (mg/100g) sample =

$$\frac{0.5 \times V_1 \times 100 \text{ ml}}{V_1 \text{ ml} \times 5 \text{ ml} \times \text{weight of the sample}} \times 100$$

Where;

V_1 = volume of oxalic acid,

3. Results and Discussion

The results are tabulated in table no. 1

Table 1: Phytochemical analysis of *Stellaria media* L.

1.	Carbohydrate (%)	17.35±0.31
2.	Protein (%)	3.30±0.16
3.	Crude Fibre (%)	13.40±0.52
4.	Ascorbic Acid (mg/100gm)	42.02±0.63
5.	Total Phenol Content (µg GAE/mg)	25.33±0.37
6.	Inhibition concentration (IC ₅₀) (µg/ml)	1021±0.67

V_2 = volume of the sample.

Total Phenol Content Estimation

The total phenol content was determined by the Folin-Ciocalteu's method [9]. 200µl of the sample extracts (1mg/ml) was taken and volume made up to 2ml. 0.3ml of Folin-Ciocalteu reagent was added. After 5mins, 0.8ml of 20% Na₂CO₃ was added and the final volume was made 5ml. Absorbance was taken by UV-Vis Spectrophotometer at 765nm after 30 minutes incubation. The amount of phenol content was determined using Gallic acid as standard. Results were expressed as µg/mg (Gallic acid equivalent/dry weight).

Antioxidant activity estimation

The antioxidant activities of the sample extracts along with standard were assessed on the basis of the radical scavenging effect of stable DPPH [10]. A solution of DPPH of concentration 0.2 mM was prepared in 70% methanol and kept overnight. Stock solution (20mg/ml) of the extract was prepared in 70% methanol. Various concentration of the extracts viz. 10, 20, 50, 100, 150, 200, 300, 400 and 500 µl were taken in different test tubes and the volume was made up to 1000µl. 1ml DPPH was added to each solution and kept at dark for 30 minutes. Ascorbic acid and Gallic acid were taken as standards. Optical density of these samples was measured at 517nm along with blank where 1ml methanol with 1 ml DPPH solution was taken. The activities of the samples are measured in terms of percent inhibition (IC₅₀) and calculated by the following formulae:

$$\text{Percent (\%)} \text{ inhibition of DPPH activity} = \frac{A-B}{A} \times 100$$


Where,

A = Optical density of the blank

B = Optical density of the sample

Statistical Analysis

The data were subjected to statistical analysis. All the assays were recorded in triplicates and the values were expressed as mean ±S.D. IC₅₀ value was calculated by plotting a graph with percent inhibition on y-axis and concentration on x-axis (Figs. 2 & 3).



*Values represented in the table are mean \pm SD of three replicates.

Fig 2: Graphics analysis of Phytochemical analysis of *Stellaria media* L.

The most common antioxidants present in herbs are vitamins C, carotenoids, flavonoids and thiol (SH) compounds, etc. There were several reports that the contribution of phenolic compounds to antioxidant activity was much greater than those of vitamin C and carotenoids [11-13]. The present investigation suggests that the major source of antioxidant capacity of *S. media* may be not from vitamin C, but rather from phenolic compounds. Moreover, these plants are also a rich source of nutritive components such as carbohydrate, crude fibre and protein that are essential for our daily diet. Therefore, the supplementation of these natural antioxidants through a balanced diet could be much more effective than the supplementation of an individual antioxidant such as vitamin C or vitamin E.

Acknowledgement

The authors are thankful to the authority of SGS Govt. P.G. College, Sidhi (M.P.) India to carried out this work.

References

- Halliwell B, Gutteridge JMC. Free radicals in biology and medicine (3rd ed.). Oxford: Oxford University Press, 1999.
- Mau JL, Chang CN, Huang SJ, Chen CC. Antioxidant properties of methanolic extracts from *Grifola frondosa*, *Morchella esculanta* and *Termitomyces albuminosus* mycelia. Food Chem, 2004; 87:111-118.
- Mondal SK, Chakraborty G, Gupta M, Mazumdar UK. Hepatoprotective activity of *Diospyros malabarica* Bark in carbon tetrachloride intoxicated rats. European Bull Drug Res, 2005; 13-25.
- Cakir A, Ahmet Yildirim MA, Duru ME, Harmandar M, Kazaz C. Isolation and Characterization of antioxidant Phenolic Compounds from the Aerial Parts of *Hypericum hyssopifolium* L. by activity-guided fractionation. J Ethnophar. 2003; 87:73-83.
- Velioglu YS, Mazza G, Gao L, Oomah BD. Antioxidant Activity and Total Phenolics in Selected Fruits, Vegetables, and Grain Products. J. Agric. Food Chem. 1998; 46:4113.
- Hedge JE, Hofreiter BT. In: Carbohydrate Chemistry 17 (Eds Whistler R L and Be Miller, J N) Academic Press New York, 1962.




Fig 3: Graphics analysis of Inhibition concentration (IC50) (mg/ml)

- Maynard AJ. (Ed): Methods in Food Analysis Academic Press New York, 1970; 176.
- Sadasivam S, Theymoli Balasubraminan. In: Practical Manual in Biochemistry, Tamil Nadu Agricultural University, Coimbatore, 1987; 14.
- Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphor-molybdc phosphotungstic acid reagents. American Journal of Enology and Viticulture. 1965; 16:144-158.
- Nooman A, Khalaf Khalaf NA, Shakya AK, Al-Othman A, El-Agbar Z, Farah H. Antioxidant activity of some common plants. Turk J. Biol. 2008; 32:51-55.
- Prior RL, Cao G, Martin A, Sofic E, McEwen J, O'Brien C, et al. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of Vaccinium species. J. Agric. Food Chem. 1998; 46:2686-2693.
- Wang H, Cao G, Prior RL. Total antioxidant capacity of fruits. J. Agric. Food Chem. 1996; 44:701-705.
- Luximon-Ramma A, Bahorun T, Crozier A. Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. J. Sci. Food Agri. 2003; 83:496-502.